



Grain irradiation and insect pests

Which insects?

A booklet from the Queensland Department of Primary Industries spells them out for Queensland:

Common name Scientific name

Rice weevil	<i>Sitophilus oryzae</i>
Lesser grain borer	<i>Rhyzoperta dominica</i>
Rust-red flour beetle	<i>Trilobium castaneum</i>
Dried fruit beetle	<i>Carpophilus dimidiatus</i>
Cadelle	<i>Tenebroides mauritanicus</i>
Bean weevil	<i>Acanthoscellides obtectus</i>
Saw-toothed grain beetle	<i>Oryzaephilus surinamensis</i>
Flat grain beetle	<i>Cryptolestes pulillus</i>
Tobacco beetle	<i>Lasioderma serricorne</i>
Tropical warehouse moth	<i>Ephestia cautella</i>
Indian meal moth	<i>Plodia interpunctella</i>
Angoumois grain moth	<i>Sitotroga cerealella</i>

Radiation sensitivity

The problem with insects is that you are dealing with more than one individual per insect. You have eggs, larvae, pupae and adults. In general the adults are the most radiation resistant.

It turns out that *Sitophilus oryzae* would need a dose of 20 krad (0.20 kGy) to kill about 99.9% of adults within 21 days. For *Oryzaephilus surinamensis* the same dose resulted in complete mortality of adults within 15 days (1). The same dose gave 'very high mortality' of adults from *Rhyzoperta dominica* (2).

Tribolium castaneum was touchier: adults died in 16 days after 50 krad (0.50 kGy) or in 12 days after 100 krad (1 kGy) (3)

Lasioderma serricorne needed 2500 Gy (2.5kGy) for 100% mortality of adults and 750 Gy or 0.75 kGy for 100% mortality of all immature stages (4).

Plodia interpunctella and *Sitotroga cerealella* are the hardest nuts to crack. All stages of these insects were given doses of about 13, 17, 25, 45 and 100 krad (0.13, 0.17, 0.25, 0.45 and 1 kGy).

It turned out that 'the life of insects treated as adults or pupae was not greatly shortened by any of the treatments.' (5).

These are only a number of the in Queensland listed insects as major storage pests. The required dose to kill adults within 1 month varies in the mentioned insects from a low 0.20 kGy to a comparatively high 2.5 kGy.

Critical points in dose range

There are two critical points in this dose range. The most obvious one is that you are not supposed to kill the grain and this can happen from 1 kGy onwards (6, 7). Nobody is interested in silos with dead, stinking grain. And the second critical point is from about 0.50 kGy onwards. Between 0.50 and 1 kGy it has been found that irradiated moulds producing aflatoxins are stimulated in producing more toxin when they get a chance to grow.

Insects love irradiated grains

It is an illusion to think that once grain has been disinfested it will stay this way. When the little critters get a chance you have them back. Well, how do they thrive on irradiated grain? It turns out they love it. So, think twice before you start irradiating.

Rhyzoperta dominica was reared on irradiated rice with doses varying from 20 to 5000 krad (0.20 to 50 kGy). This resulted in an increase in adult life-span and fecundity of the next generation (8).

Tribolium castaneum had similar preferences. Reared on irradiated diets the numbers in offspring of the second, third and fourth generation were significantly higher than the offspring on unirradiated diet (9).

References

1. Hoedaya, M.S. et al. 1973. Radiation effects on four species of insects etc. 281-294. Pasar Jum'at Research Centre, National Atomic Energy Agency, Jakarta, Indonesia.
2. Abdul Matin, A.S. et al. 1974. Susceptibility of *Rhyzopertha dominica* to ionizing radiation. *Journal of Stored Products research* 10(3/4) 199-207. - *Review of Applied Entomology -Series A* 1975 vol.63 No.10 p.1052 no.3864.
3. Adem, E. et al. 1979. Responses of *Prostephanus* etc. and *Tribolium castaneum* to gamma irradiation. *Canadian Entomologist* 111 (10) 1111-1114. - *Review of Applied Entomology - Series A* 1980 vol. 68 NO.6- p.392 no.3124
4. Els, J. M. et al. 1978, recd. 1980. The lethal effect of gamma irradiation on *Lasioderma srricorne*. *Phytophylactica* 10 (4) 127-128. Tobacco Research Institute , Private Bag, Rustenburg, 0300 South Africa. - *Review of Applied Entomology - Series A* 1980 vol.68 No.12 - p.826 no 6652.
5. Cogburn, R.R. et al. 1966. Gross effects of gamma radiation on the Indian-meal moth and the Angoumois grain moth. *J. econ. Ent.* 59 no.3 pp. 682-685. - *Review of Applied Entomology* vol.54. - p.495
6. Pesson, P. 1963. Utilisation des radiations ionisantes pour la protection des denrees contre les insects nuisibles. *Industr. agric.* 80 no.3 pp. 211-225.- *Review of Applied Entomology* vol.54, 1966. P 58.
7. Wills, P.A. 1988. Critical review of a submission by H. Julius , Food Irradiation and Moulds: a time bomb. An ANSTO report p.34.
8. Wiendl, F. et al. 1974, recd. 1976. Mortalidade e reproducao de *Rhyzopertha dominica*. *Anais da Sociedade Entomologica do Brasil.* 3 (1) p. 34-43. - *Review of Applied Entomology - Series A* 1977. Vol. 65 No.2, no.875
9. Tilton, E.W. et al. 1973. Progeny production by *S. oryzae* and *Tribolium castaneum* reared for several generations on irradiated diets. *Journal of the Georgia Entomological Society* 8 (3) pp. 168-173. - *Review of Applied Entomology - Series A* 1974 vol. 62 No.11. p.1222 no. 4603.

H. Julius

1999 Friends of the Earth

feel free to copy